

KLICKMER™ T CELL IMMUNE MONITORING AND BEYOND!

**YOUR DEXTRAMER® WILL BE READY TO GO
IN JUST ONE KLICK**

Add biotinylated protein to the fluorescent Klickmer™ and create your personalized Dextramer® reagent. Push the power of Dextramer® beyond the detection of epitope specific T cells.

KLICKMER™ – DESIGN YOUR OWN DEXTRAMER® MOLECULE

WHAT IS KLICKMER™?

A Klickmer™ is a Dextramer® backbone carrying multiple acceptor sites, enabling binding of a high number of biotinylated molecules. To generate high avidity multimers displaying the molecule of choice. Klickmer™ reagents carry PE/FITC/APC label, and an unlabeled version is also available.

GET INSPIRED BY KLICKMER™...

Klickmer™ represents the scaffold to which potentially any biotinylated molecule can be attached. This multi-purpose tool serves a range of applications and can help you address your scientific questions.

APPLICATIONS

- Highly sensitive detection of antigen-specific T cells and B cells.
- Simulation of antigen presentation to B or T cells
- Identification of ligands for low-affinity interactions or orphan receptors.
- Amplification of the signal on your cell populations by flow cytometry, or other fluorescence assays.
- Isolation of specific antigen-presenting cells.

REFERENCE

- 1) Dolton, G, et al. Comparison of peptide-major histocompatibility complex tetramers and dextramers for the identification of antigen-specific T cells. Clin Exp Immunol. 2014; 177[1]:47-63.
- 2) Rius, C, et al. Peptide-MHC Class I Tetramers Can Fail To Detect Relevant Functional T Cell Clonotypes and Underestimate Antigen-Reactive T Cell Populations. J Immunol. 2018; 200 [7]: 2263-2279.
- 3) Dolton, G, et al. Optimized Peptide-MHC Multimer Protocols for Detection and Isolation of Autoimmune T-Cells. Front Immunol. 2018;9:1378.
- 4) Schøller, J, et al. A recombinant human HLA-class I antigen linked to dextran elicits innate and adaptive immune responses. J Immunol Methods. 2010; 360[1-2]: 1-9.
- 5) Mörner, A, et al. Immunization with recombinant HLA classes I and II, HIV-1 gp140, and SIV p27 elicits protection against heterologous SHIV infection in rhesus macaques. J Virol. 2011;85[13]:6442-52.
- 6) Gui-Bo, Y, et al. Immunization with recombinant macaque major histocompatibility complex class I and II and human immunodeficiency virus gp140 inhibits simian-human immunodeficiency virus infection in macaques. J Gen Virol. 2012;93(Pt 7):1506-18.
- 7) Chancellor, A, et al. CD1b-restricted GEM T cell responses are modulated by Mycobacterium tuberculosis mycolic acid meromycolate chains. Proc Natl Acad Sci U S A. 2017;114[51].
- 8) Luque, S, et al. A multicolour HLA-specific B-cell FluoroSpot assay to functionally track circulating HLA-specific memory B cells. J Immunol Methods. 2018;462:23-33.
- 9) Kasmar AG, et al. Cutting Edge: CD1a tetramers and dextramers identify human lipopeptide-specific T cells ex vivo. J Immunol. 2013;191[9]:4499-503.
- 10) Lolli F, et al. Increased CD8+ T cell responses to apoptotic T cell-associated antigens in multiple sclerosis. J Neuroinflammation. 2013;10:94.
- 11) Schøller J, et al. A recombinant human HLA-class I antigen linked to dextran elicits innate and adaptive immune responses. J Immunol Methods. 2010;360[1-2]:1-9.

KLICKMER™

