

Recommended use U-Load Dextramer® is a fluorescent dextramer backbone optimized for attachment of MHC-peptide monomers. U-Load Dextramer® is used in combination with easYmer® or U-Load MHC-II to generate U-Load Dextramer® MHC reagents that can be used for detection of antigen-specific T cells using flow cytometry (PE, APC, FITC).

For research use only. Not for use in diagnostic or therapeutic procedures.

Reagents provided

- U-Load Dextramer®: Fluorescent (PE, APC, FITC) dextramer backbone or without fluorochrome (NONE)
- U-Load Dextramer® Dilution buffer: PBS, 0.5% BSA, 7.5 mM NaN₃, pH 7

Sizes

Tests	Content	
	U-Load Dextramer®	U-Load Dextramer® Dilution Buffer
20	1 vial (40 µL)	1 vial (2 mL)
50	1 vial (100 µL)	1 vial (2 mL)
150	1 vial (300 µL)	1 vial (2 mL)
500	1 vial (1000 µL)	2 vials (2 mL)

Storage Store in the dark at 2-8°C.

Precautions Contains sodium azide (NaN₃), a chemical highly toxic in pure form. At product concentrations, though not classified as hazardous, sodium azide may react with lead and copper, plumbing to form highly explosive build-ups of metal azides. Upon disposal, flush with large volumes of water to prevent metal azide build-up in plumbing. As with any product derived from biological sources, proper handling procedures should be used. For professional users.

Recommended protocols MHC I: See "Protocol for Preparation of MHC I-peptide monomer and fluorescent U-Load Dextramer® MHC I" and "General staining procedure MHC Dextramer – PBMC's" (www.immudex.com/resources/protocols/).

MHC II: See "Protocol for Preparation of MHC II-peptide monomer and fluorescent U-Load Dextramer® MHC II" and "General staining procedure MHC Dextramer – PBMC's" (www.immudex.com/resources/protocols/).

Symbols See www.immudex.com/symbols

Technical support E-mail: customer@immudex.com

Telephone: +45 3110 9292 (Denmark)

Manufacturer Immudex, Bredevej 2A, DK-2830 Virum, Denmark