

<b>Recommended use</b>	<p>TCR Dextramer® reagent is recommended for identification and enumeration of specific antigen-presenting cells in cell samples using flow cytometry.</p> <p>For research use only. Not for use in diagnostic or therapeutic procedures.</p>
<b>Reagents provided</b>	<p>TCR Dextramer® reagent is uniquely identified by its account product number and fluorophore e.g., DTCRU0035 / PE</p> <p>TCR Dextramer® reagent consists of a fluorescent (PE, APC, FITC) dextran backbone, displaying multiple TCR monomers.</p> <p>TCR Dextramer® reagent is also available without fluorochrome (NONE).</p> <p>TCR Dextramer® reagent is provided at a concentration of 32nM in buffer containing 1% bovine serum albumin (BSA) and 15 mM NaN<sub>3</sub>, pH 7.2.</p> <p>This is a custom-made product based on TCR alpha and beta chain sequences provided by customer.</p>
<b>Sizes</b>	See test size on label
<b>Storage</b>	Store in the dark at 2-8°C.
<b>Precautions</b>	<p>Contains sodium azide (NaN<sub>3</sub>), a chemical highly toxic in pure form. At product concentrations, though not classified as hazardous, sodium azide may react with lead and copper, plumbing to form highly explosive build-ups of metal azides. Upon disposal, flush with large volumes of water to prevent metal azide build-up in plumbing.</p> <p>As with any product derived from biological sources, proper handling procedures should be used.</p> <p>For professional users.</p>
<b>Symbols</b>	See <a href="http://www.immudex.com/symbols">www.immudex.com/symbols</a>
<b>Technical support</b>	<p>E-mail: <a href="mailto:customer@immudex.com">customer@immudex.com</a></p> <p>Telephone: +45 3110 9292 (Denmark). +1 (703) 766 4688 (US)</p>
<b>Manufacturer</b>	Immudex, Bredevej 2A, DK-2830 Virum, Denmark