

Products	<p>CSS009_x TCR Dextramer® reagent is uniquely identified by its cat. no. and fluorophore e.g., CSS009_x / PE.</p> <p>TCR Dextramer® reagent consists of a fluorescent (PE) dextran backbone, displaying an optimal number of TCR Monomers. The TCR specificity and affinity (if applicable) are mentioned on the label, e.g., HLA-A*0102 / SLLMWITQV and 1 nM.</p> <p>CSS008_4 TCR Dextramer® reagent is uniquely identified by its account product number and fluorophore, e.g., DTCRU0035 / PE.</p> <p>TCR Dextramer® reagent consists of a fluorescent (PE, APC, FITC) dextran backbone, displaying multiple TCR monomers. TCR Dextramer® reagent is also available without fluorochrome (NONE).</p> <p>This is a custom-made product based on TCR alpha and beta chain sequences provided by the customer.</p>
Recommended use	<p>TCR Dextramer® reagent is recommended for identification and enumeration of specific antigen-presenting on cells in cell samples using flow cytometry.</p> <p>For research use only. Not for use in diagnostic or therapeutic procedures.</p>
Concentration	<p>TCR Dextramer® reagent is provided at a concentration of 32 nM in 50 mM Tris-HCl buffer containing 1% bovine serum albumin (BSA) and 15 mM NaN₃, pH 7.2.</p>
Sizes	<p>See test size on label.</p>
Storage	<p>Store in the dark at 2-8°C.</p>
Precautions	<p>Contains sodium azide (NaN₃), a chemical highly toxic in pure form. At product concentrations, though not classified as hazardous, sodium azide may react with lead and copper, plumbing to form highly explosive build-ups of metal azides. Upon disposal, flush with large volumes of water to prevent metal azide build-up in plumbing.</p> <p>As with any product derived from biological sources, proper handling procedures should be used.</p> <p>For professional users.</p>
Symbols	<p>See www.immudex.com/symbols</p>
Technical support	<p>E-mail: customer@immudex.com</p> <p>Telephone: +45 3110 9292 (Denmark). +1 (703) 766 4688 (US)</p>
Manufacturer	<p>Immudex, Bredevej 2A, DK-2830 Virum, Denmark</p>